

Important Questions for Class 9

Maths

Chapter 12 – Statistics

Section A

1. If the mean of 2,4,6,8,x,y **is 5 then find the value of x+y**.

Ans: 10

2. Write the class mark of 90–110 group.

Ans: 100

3. If the ratio of mean and median of a certain data is 2: 3, then find the ratio of its mode and mean.

Ans: 5:2

4. Tally marks are used to find

Ans: Frequency

5. The following marks were obtained by the students in a test. 81,72,90,90,86,85,92,70,71,83,89,95,85,79,62. What is the range?

Ans: 33

6. In a histogram, each class rectangle is constructed with base as

Class IX Maths



1. frequency

2. class interval

3. range

4. size of the class

Ans: class interval

Section B

7. The mean of 10 numbers is 20. If 5 is subtracted from every number, what will be the new mean?

Ans: Let x_1, x_2, \dots, x_{10} be 10 numbers with their mean equal to 20.

Then,

$$\overline{X} = \frac{1}{n} \left(\sum x_i \right) 20 = \frac{x_1 + x_2 + \dots + x_{20}}{10}$$

 $\Rightarrow 200 = x_1 + x_2 + \dots + x_{10}$ New numbers are $x_1 - 5, x_2 - 5, \dots, x_{10} - 5$

Let \overline{X} be the mean of new number.

$$\overline{X} = \frac{(x_1 - 5) + (x_2 - 5) + \dots + (x_{10} - 5)}{10} \quad X = \frac{(x_1 + x_2 + \dots + x_{10}) - 5 \times 10}{10}$$
$$= \frac{200 - 50}{10} = 15$$

8. Find the mean of the first 10 even natural no.

Ans:
$$\bar{x} = \frac{2+4+6+8+10+12+14+16+18+20}{10} = \frac{10\times11}{10} = 11$$

9. Calculate the mean for the following distribution.

Class IX Maths



 x
 5
 6
 7
 8
 9

 f
 4
 8
 14
 11
 3

Ans: 7.025

10. Find the median of 37, 31, 42, 43, 46, 25, 39, 45, 32.

Ans: 39

11. Find the mode of the following series. 25, 23, 22, 22, 24, 27, 27, 25, 23, 22, 26, 32.

Ans: 22

12. If the median of a series of data is 3 and mean is 2 then, find the mode.

Ans: 5

Section C

13. Find the median of the following data 19, 25, 59, 48, 35, 31, 30, 32, 51. If 25 is replaced by 52, what will be the new median.

Ans: 32, 35

14. If the mean of the following distribution is 6, then find the value of p.

x 2 4 6 10 p+5**f** 3 2 3 1 2

Ans: 7



15. If the mean of five observations x, x+2, x+4, x+6, x+8 is 11, find the mean of first three observation.

Ans: 9

16. The mean of 5 numbers is 18. If one number is excluded, their mean is 16, find the excluded number.

Ans: 26

17. The following observations have been arranged in ascending order. If the median of the data is 63, find the value of x.

29, 32, 48, 50, *x*, *x* + **2, 72**, 78, 84, 95

Ans: 62

Section D

18. Find the value of x and y in the following distribution if it is known that the mean of the distribution is 1.46.

Ans: x = 76, y = 38

19. The mean monthly salary of 10 members of a group is Rs 1445, one more member whose monthly salary is Rs 1500 has joined the group. Find the mean monthly salary of 11 members of the group.

Ans: Rs 1450

20. Given below is a cumulative frequency distribution table showing the age of people living in a locality.



Age in years	Number of persons
Above 108	0
Above 96	1
Above 84	3
Above 72	5
Above 60	20
Above 48	158
Above 36	427
Above 24	809
Above 12	1026
Above 0	1124

Prepare a frequency distribution table.

Ans:

Marks	0-12	12 - 24	24 - 36	36-48	48 - 60	60 - 72	72 - 84	<u>84–96</u>	96-108
Person	98	217	382	269	138	15	2	2	1

21. If $\mathbf{x}, \mathbf{x}_2, \dots, \mathbf{x}_n$ are n



values of a variable x such that $\sum_{i=1}^{n} (x_i - 2) = 110$ and $\sum_{i=1}^{n} (x_i - 5) = 20$ find the value of n and mean.

Ans:
$$\sum_{i=1}^{n} (x_i - 2) = 110$$

 $\Rightarrow \sum_{i=1}^{n} x_i - 2n = 110$
 $\Rightarrow \sum_{i=1}^{n} x_i = 2n + 110$
 $\sum_{i=1}^{n} (x_i - 5) = 20$
 $\Rightarrow \sum_{i=1}^{n} x_i - 5n = 20$
 $\Rightarrow \sum_{i=1}^{n} x_i = 5n + 20$
 $\Rightarrow 5n + 20 = 2n + 110$
 $\Rightarrow 3n = 90$
 $\Rightarrow n = 30$
 $\Rightarrow \overline{x} = \frac{\sum_{i=1}^{n} x_i}{n} = \frac{2n + 110}{n} = \frac{2(30) + 110}{30} = \frac{170}{30} = \frac{17}{3}$
 $n = 30, mean = \frac{17}{3}$

22. The mean of 200 items was 50. Later on, it was discovered that the two items were misread as 92 and 8 instead of 192 and 88. Find the correct mean.

Ans:



$$\overline{x} = \frac{\sum_{i=1}^{n} x_i}{n}$$
$$\implies 50 = \frac{\sum_{i=1}^{n} x_i}{200}$$

 $\Rightarrow \sum_{i=1}^{n} x_i = 200 \times 50 = 10000$

For new mean,

$$\Rightarrow \sum_{i=1}^{n} x_i = 10000 - (92 + 8) + (192 + 88) = 10180$$
$$\Rightarrow \overline{x} = \frac{\sum_{i=1}^{n} x_i}{n} = \frac{10180}{200} = 50.9$$

23. Find the value of p, if the mean of the following distribution is 20.

Х	15	17	19	20+p	23
frequency	2	3	4	5p	6

Ans: 1

Important Questions

1 Marks questions

1. If the mean of 2,4,6,8,x, y is 5 then find the value of x+y.

Ans: 10

2. Write the class mark of 90–110 group.

Ans: 100



3. If the ratio of mean and median of a certain data is **2**: **3**, then find the ratio of itsmode and mean.

Ans: 5:2

4. Tally marks are used to find

Ans: Frequency

5. The following marks were obtained by the students in a test. 81,72,90,90,86,85,92,70,71,83,89,95,85,79,62. What is the range?

Ans: 33

6. In a histogram, each class rectangle is constructed with base as

- 1. frequency
- 2. class interval
- 3. range
- 4. size of the class

Ans: class interval

7. Find the range of the following data: 25, 20, 30, 18, 16, 15.

- (a) 15
- (b) **10**
- (c) 5
- (d) 20

Ans: (a) 15

Class IX Maths



8. Find the median of the given data: 7, 8, 7, 7, 9, 10, 13.

- (a) 7
- (b) 9
- (c) **8**
- (d) 10
- **Ans:** (c) 8

9. Find the mode of the given data: 7, 9, 11, 13, 9, 13, 9, 9, 7, 8.

- (a) 10
- (b) **9**
- (c) 11
- (**d**) 8
- **Ans:** (b) 9

10. Find the mean of the first five multiples of 3?

- (a) 9
- (b) **12**
- (c) 14
- (d) None of these
- **Ans:** (a) 9

11. The mean of 7 observations is 20. If the mean of the first 4 observations is 12 & that of last 4 observations is 28, find the 4th observation ?



Ans: Since mean of 7 observations = 20

 \therefore Total of 7 observations = $20 \times 7 = 140$

:. Mean of first 4 observations $=12 \implies$ Total of first 4 observations $=12 \times 4 = 48$

 \therefore Mean of last 4 observations = 28 \Rightarrow Total of first 4 observations = 4×28=92

 \therefore Total of 7 observations +4th observation = 48+92

 $140 + 4^{\text{th}}$ observation = 140

 $\Rightarrow 4^{\text{th}} \text{ observation } = 140 - 140 = 0$

12. What is the upper limit of the interval: 20–23?

- (a) 20
- (b) 23
- (c) 22
- (d) None of these

Ans: (b) 23

13. What is class size of interval 10, 12, 14, 16, 18 ?

- (a) 2
- **(b)** 1
- (c) 10
- (d) 18

Ans: (a) 2

14. Find the class mark of the interval 15.7–25.7?



(a) 15.3

- (b) 16.3
- (c) 17.3
- (d) 20.7
- **Ans:** (a) 20.7

15. what is the mid-point of class interval 12.3–22.3?

- (a) 17.3
- (b) 15.3
- (c) 18.3
- (d) 16.7

Ans: (a) 17.3

16. The median of the following observations arranged in ascending order is 40. find x

15, 12, 11, 14, x+2, x+4, 32, 30, 41, 35

Ans: No. of observation =10 which is even

:. Median is the average of $\left(\frac{n}{2}\right)^{th}$ and $\left(\frac{n}{2}+1\right)^{th}$ observations

So, 5th observation = x + 2 and 6th observation = x + 4

Median
$$=\frac{(x+2)+(x+4)}{2}$$

$$40 = \frac{2x+6}{2}$$



40 = x + 3

 $\therefore x = 37$

17. What is the class Mark of the interval 15-20?

(A) 15

(B) 20

(C) **17.5**

(D) none of these

Ans: (C) 17.5

18. What is the range of interval 15-20?

- (A) **5**
- **(B) 10**
- (C) 15

(D) none of these

Ans: (A) 5

19. What is the class - size of the interval 15-20?

(A) **5**

(B) 10

(C) 15

(D) none of these

Ans: (A) 5

Class IX Maths



20. Find out the mean of following data. 5,10,15,20,25,30.

- (A) **16.5**
- **(B) 17.5**
- (C) **18.5**
- **(D)** none of these
- **Ans:** (B) 17.5

21. Find the arithmetic mean of first 6 natural numbers?

- (a) **3.5**
- (b) **4.5**
- (c) 2.5
- (d) none of these
- **Ans:** (a) 3.5

22. What is the mid-point of interval 3-6 ?

- (a) **3.5**
- (b) **4.5**
- (c) 5.5
- (d) none of these
- **Ans:** (b) 4.5

23. Find out the range of the following: 5, 10, 15, 20, 25, 30.

(a) 25

Class IX Maths



(b) **20**

(c) **30**

(d) none of these

Ans: (a) 25

24. Find out the mode of the following: 5, 4, 3, 5, 6, 6, 6, 5, 4, 5, 5, 3, 2, 1.

- (a) **6**
- (b) 4
- (c) **5**
- (d) none of these

Ans: (c) 5

25. What is the class size of the intervals 10-20 ?

- (a) 10
- (b) **5**
- (c) 15
- (d) 20
- **Ans:** (a) 10

26. What is the upper class limit of the class 37-43?

- (a) **37**
- (b) **40**
- (c) **43**



(d) none of these

Ans: (c) 43

27. What is the lower class limit of the class 37-43?

(a) **37**

- (b) **40**
- (c) **43**
- (d) none of these

Ans: (a) 37

28. Find the median of the following data: 15, 35, 18, 26, 19, 25, 29, 20, 27, 30.

- (a) 25.5
- (b) 24.5
- (c) 26.5
- (d) none of these

Ans: (a) 25.5

2 Marks **Questions**

1. The mean of 10 numbers is 20, If 5 is subtracted from every number, what will be the new mean.

Ans: 15

2. Find the mean of first 10 even natural no.



Ans: 11

3. Calculate the mean for the following distribution.

x 5 6 7 8 9 f 4 8 14 11 3

Ans: 7.025

4. Find the median of 37, 31, 42, 43, 46, 25, 39, 45, 32.

Ans: 39

5. Find the mode of following series. 25, 23, 22, 22, 24, 27, 27, 25, 23, 22, 26, 32. Ans: 22

6. If the median of a series of data is 3 and mean is 2 then, find the mode.

Ans: 5

7. If the mean of 5 observation x, x+4, x+8, x+12, x+16 is 13, find the mean of the observations?

Ans:
$$\overline{x} = \frac{\sum x_i}{n}$$

$$\Rightarrow 13 = \frac{x + (x+4) + (x+8) + (x+12) + (x+16)}{5}$$

$$\Rightarrow 5 \times 13 = 5x + 40$$

 $\Rightarrow 13 = x + 8$



 $\therefore x = 5$

 \therefore The five given observations are 5, 9, 13, 17, 21

$$\overline{x} = \frac{5+9++13+17+21}{5} = 12.8$$

8. The class marks of the observations are 17, 21, 25, 29, 33, 37, 41, 45. Find the class intervals.

Ans: Class marks are 17, 21, 25, 29, 33, 37, 41 and 45

Class size = 21 - 17 = 25 - 21 = 4

Half of class size = $\frac{4}{2} = 2$

Thus, Class intervals are:

17-2=15	&	17+2=19	i.e	15-19
21-2=19	&	21+2=23	i.e	19-23
25-2=23	&	25+2=27	i.e	23-27
29-2=27	&	29+2=31	i.e	27-31
33-2= <mark>31</mark>	&	33+2=35	i.e	31-35
37-2=35	&	37+2=39	i.e	35-39
41-2=39	&	41+2=43	i.e	39-43
45-2=43	&	45+2=47	i.e	43-47



9. The value of π up to 15 decimal places is: \$3.419078023195679\$

(i) List the digits from 0 to 9 & make frequency distributions of the digit after the decimal points.

(ii) What are the most and the least frequently occurring digits?

Digits	Tally Marks	Frequency
0	Ш	2
1	П	2
2	I	1
3		1
4	I	1
5		1
6		1
7		2
8		1
9		3

Ans: (i) Frequency distribution table

(ii) Most frequency occurring digits =9

Least frequently occurring digits = 2, 3, 4, 5, 6, 8



10. A random survey of the number of children of various age grout playing in the park was found:

Age[inyears]1-22-33.55-7 $7 \cdot 10$ No. of children3571013

Draw a histogram to represent the data above?

Ans: we will calculate the adjusted frequencies [AF] for histogram because the class intervals are not of equal width.

Minimum class size [CS] = 1

Age (In Years)	Frequency	Class Size(CS)	$AF = \frac{Minimum CS}{CS \text{ of this class}} \times \text{ Its frequency}$
1-2	3	1	$\frac{1}{1} \times 3 = 3$
2-3	5	1	$\frac{1}{1} \times 5 = 5$
3-5	7	2	$\frac{1}{2} \times 7 = 3.5$
5-7	10	2	$\frac{1}{2} \times 10 = 5$
7-10	13	3	$\frac{1}{3} \times 13 = 4.3$



To get the required histogram, draw rectangles with base equal to class interval and heights equal to the corresponding adjusted frequencies.



11. The mean of 20, 8, 12, 13, 15, P is **30**. Find the value of P?

Ans. Mean $= \overline{x} = \frac{20 + 8 + 12 + 13 + 15 + P}{6}$ $30 = \frac{68 + P}{6}$ $\Rightarrow 180 = 68 + P$ $\therefore P = 180 - 68 = 112$

12. Find the mode of the following data: 14, 25, 14, 14, 25, 24, 20, 28, 18, 20.

Ans: Arrange the given data in ascending order:

Class IX Maths



14, 14, 14, 18, 20, 20, 24, 25, 25, 28

The value 14 occurs most frequently i.e. 3 times as observed.

So, Mode is 14

13. Find the median of 5, 7, 10, 9, 5, 12, 15, 12, 18, 20. If 9 is replaced by 14, what will be the new median?

Ans: Arrange the given data in ascending order:

5, <mark>5, 7, 9, 10, 12</mark>, 15, 18, 20

Here, n = 10 [even number]

So, Median = $\frac{5^{\text{th}} \text{ observation } + 6^{\text{th}} \text{ observation}}{2} = \frac{10+12}{2} = 11$

After replacing 9 by 14:

5, **5**, **7**, **10**, **12**, **12**, **14**, **15**, **18**, **20**

Now, 5^{th} observation = 6^{th} observation = 12

So, New Median $=\frac{12+12}{2}=12$

14. The average mark of boys in an examination is 68 & that of girls in 89. If the average mark of all candidates in that examination is 80, find the ratio of the no. of boys to the number of girls that appeared in the examinations.

Ans: Let number of boys be x & girls be y.

 \therefore Total marks of boys = $68 \times x = 68x$

 \therefore Total marks of girls = 89 × y = 89 y

Hence, total marks for boys & girls = 68x + 89y

Also, total of boys & girls = x + y & average for all the candidates = 80



 $\Rightarrow 80 = \frac{68x + 89y}{x + y}$ 80x + 80y = 68x + 89y 80x - 68x = 89y - 80y 12x = 9y

- $\Rightarrow \frac{x}{y} = \frac{9}{12} = \frac{3}{4}$
- \therefore Ratio of boys & girls = 3:4

15. The mean of 6 numbers is 30. If one number is excluded, their mean is 24 . Find the excluded number.

Ans: Here, n = 6, $\overline{x} = 30$

$$\therefore \ \overline{x} = \frac{1}{n} \left(\sum x_i \right) \Longrightarrow \sum x_i = n \overline{x}$$

$$\sum x_i = 6 \times 30 = 180$$

So, total of 6 numbers =180

Let the excluded number be a.

Now, total of 5 number is 180 -a

Mean of 5 number = $\frac{180-a}{5}$

 $\Rightarrow 24 = \frac{180 - a}{5}$

 $\Rightarrow 24 \times 5 = 180 - a$

 $\Rightarrow a = 180 - 120$

$$\Rightarrow a = 60$$



Thus, excluded number is 60.

16. The median of the observation 11, 12, 14, 18, x+2, x+4, 30, 32, 35, 41, arranged in ascending order is 24. find the value of x.

Ans: Number of observation, n = 10

Since n is even,

 $\therefore \text{ median} = \frac{\left[\frac{n}{2}\right]^{th} \text{ observation } + \left[\frac{n}{2} + 1\right]^{th} \text{ observation}}{2}$

$$\Rightarrow 24 = \frac{5^{th} \text{ observation}}{2} + 6^{th} \text{ observation}$$

$$\Rightarrow 24 = \frac{(x+2) + (x+4)}{2}$$

$$\Rightarrow 24 = \frac{2x+6}{2} \Rightarrow 24 = x+3$$

 $\Rightarrow x = 21$

So, x = 21

17. Find the median of the following data: 25, 34, 31, 23, 22, 26, 35, 28, 20, 32.

Ans: Arrange the data in ascending order:

20, 22, 23, 25, 26, 28, 31, 32, 34, 35.

Hence, the no. of observation n = 10 (even)

$$\Rightarrow$$
 median = $\frac{5^{th}$ observation + 6^{th} observation 2

$$=\frac{26+28}{2}=27$$

Class IX Maths



So, median of the data is 27.

18. In x standard, these are three section A, B, C with 25, 40 and 30 students respectively. The average mark of section A is 70%, of section B is 65% and of section B is 50%. Find the average marks of the entire X standard.

Ans: $n_1 = 25, n_2 = 40, n_3 = 35$

 $\bar{x}_1 = 70, \bar{x}_2 = 65$, and $\bar{x}_3 = 50$

Let \overline{X} , denote the average mark of the entire X standard.

Then,
$$\bar{X} = \frac{n_1 X_1 + n_2 X_2 + n_3 X_3}{n_2 + n_2 + n_3}$$

= $\frac{25 \times 70 + 40 \times 65 + 35 \times 50}{25 + 40 + 35}$
= $\frac{1750 + 2600 + 1750}{100}$
= $\frac{6100}{100} = 61$

So, the average marks of the entire X standard is 61%.

19. If \overline{x} is a mean of $x_1, x_2, x_3 - - - - x_n$ then the mean of $ax_1, ax_2, - - - ax_n$ is $a\overline{x}$ where 'a' is any number different from 2 era i.e. if each observation is multiplied by a non 2 era number a, then the mean is also multiplied by a.

Ans: We have:
$$\overline{X} = \frac{1}{n} \left[\sum_{i=1}^{n} X_i \right]$$

Let X be the mean of $ax_1, ax_2...ax_n$. Then,

$$X = \frac{ax_1 + ax_2 + \ldots + ax_n}{n}$$



$$= \frac{a(\mathbf{x}_1 + \mathbf{x}_7 + \dots + \mathbf{x}_n)}{n}$$
$$= a\left[\frac{1}{n}\sum_{i=1}^n X_i\right]$$

 $= a\overline{X}$

Thus, mean gets multiplied by a.

20. The class marks of a distribution are 26, 31, 41, 36, 46, 51, 56, 61, 66, 71. Find the true class limits.

Ans: Class size is the difference between any two consecutive class marks because class marks are uniformly spaced.

 \therefore Class size = 31-26=5

If a is the class mark of a class interval of size h, then the lower and upper limits of the class intervals are $a - \frac{h}{2}$ and $a + \frac{n}{2}$ respectively.

Here h = 5

:. Lower limit of first class interval = $26 - \frac{5}{2} = 23.5$ and upper limit of first class interval = $26 + \frac{5}{2} = 28.5$

∴ First class interval is 23.5-28.5

So, true class limits:

23.5-28.5, 28.5-33.5, 33.5-38.5, 38.5-43.5, 43.5-48.5, 48.5-53.5

21. The marks obtained by 15 students in an examination are given below;

125, 130, 130, 120, 141, 146, 162, 163, 169, 173, 179, 188, 192, 195, 199.

Form a cumulative frequency table with class interval of length 20.

Class IX Maths



Ans:

Class Interval	Tally Marks	Frequency	Cumulative frequency
120-140		4	4
140-160	=	2	6
160-180	Ш	5	11
180-200		4	15
Total		15	15

22. For the following data, draw a histogram and a frequency polygon.

Mark	0-10	10-20	20-30	30-40	40-50
No. of Students	5	10	4	6	7

Ans:





23. If X is the mean of n observation x_1, x_2, \ldots, x_n , then prove that the mean of $x_1 - a, x_2 - a, \ldots, x_n - a$, is X - a, where a is any real number.

Ans: We have $\overline{X} = \frac{1}{n} \left[\sum_{i=1}^{n} X_{i} \right]$

Let X be the mean of $x_1 - a_2, x_2 - a_3, \dots, x_3 - a$.

Then,
$$X = \frac{(x_1 - a) + (x_2 - a) + \dots + (x_n - a)}{n}$$

$$=\frac{\left(x_1+x_1+\ldots,x_n\right)-na}{n}$$

$$=\frac{1}{n}\left[\sum_{i=1}^{n}X_{i}\right]-\frac{na}{a}$$

$$=\overline{X}-a$$

Class IX Maths



Hence, proved.

24. The mean of 16 numbers is 8. If 2 is added to every number, what will be new mean?

Ans: Let $x_1, x_2x_3, \ldots, x_{16}$ be 16 numbers with their mean equal to 8.

Then,
$$\overline{X} = \frac{1}{n} \left(\sum x_i \right)$$

$$\Rightarrow 8 = \frac{x_1 + x_2 + \dots + x_{15}}{16}$$

$$\Rightarrow 16 \times 8 = x_1 + x_2 + \dots + x_{16}$$

New numbers are : $x_1 + 2, x_2 + 2, x_3 + 2, \dots, x_{16} + 2$

Let *X* Be the mean of new numbers. Then,

$$X = \frac{(x_1 + 2) + (x_2 + 2) + \dots + (x_{16} + 2)}{16}$$
$$= \frac{(x_1 + x_2 + \dots + x_{16}) + 2 \times 16}{16}$$
$$= \frac{128 + 32}{16}$$
$$= \frac{160}{16} = 10$$

25. Calculate the mean from the given data.

Mark	15	20	25	30	35	40
No. of students	6	7	12	14	15	6

Ans:





26. The following table gives the mark scored by 50 students in an entrance examination:

Mark	0-20	20-40	40-60	60-80	80-100
No. of students	7	6	13	16	8



From this table find:

(i) the less than series and

(ii) the more than series.

Ans: (i) Less than cumulative frequency table.

Mark obtained	Number of students (Cumulative frequency)
Less than 20	7
Less than 40	13
Less than 60	26
Less than 80	42
Less than 100	50

(ii) More than cumulative frequency table.

Mark obtained	Number of students (Cumulative frequency)
More than 0	50
More than 20	43
More than 40	37
More than 60	24
More than 80	8



More than 100

0

27. Find the sum of the deviations of the various values 3, 4, 6, 8, 14 from their mean.

Ans: Recall that the deviations of the values $x_1, x_2, x_3, \dots, x_n$ about A are

 $X_1 - A, x_2 - A, x_3 - A, \dots, x_n - A$

Let *x* be the deviations of the values 3, 4, 6, 8, 14. Then,

$$X = \frac{3+4+6+8+14}{5} = \frac{35}{5} = 7$$

Now, sum of the deviations of the values 3, 4, 6, 8 and 14 , From their mean $\overline{X} = 7$ is given by

(3-7)+(4-7)+(6-7)+(8-7)+(14-4) = -4-3-1+1+7=0

28. The mean of 40 observations was 200. It was detected on rechecking that the value of 65 was wrongly copied as 25 for computation of mean. Find the correct mean.

So,
$$X = \frac{1}{n} \left(\sum x \right) \Rightarrow 200 = \frac{1}{40} \left(\sum x_i \right)$$

$$\sum x_1 = 200 \times 40 = 8000$$

Ans: $n = 40, \ \bar{X} = 200$

:. Incorrect value of $\sum x = 8000$ correct value of $\sum x_i$ = incorrect value of $\sum x_1$ - incorrect item + correct item = 8000 - 25 + 65 = 8040

 $\therefore \text{ correct mean } = \frac{8040}{40} = 201$



29. It \bar{X} is the mean of n observation x_1, x_2, \dots, x_2 , then prove that $\sum_{i=1}^{n} (X_i - \bar{X}) = 0$

i.e., the algebraic sum of deviations from mean is zero.

Ans: We have
$$\overline{X} = \frac{1}{n} \left[\sum_{i=1}^{n} x_i \right]$$

 $\Rightarrow n\overline{X} = \sum_{i=1}^{n} x_i$
Now, $\sum_{i=1}^{n} (x_i - \overline{X}) = (x_i - \overline{X}) + (x_2 - \overline{X}) + \dots + (x_n - \overline{X})$
 $= (x_1 + x_2 + \dots + x_n) - n\overline{X}$
 $= \sum_{i=1}^{n} x_i - n\overline{X}$
 $= n\overline{X} - n\overline{X}$
 $= 0$
So, $\sum_{i=1}^{n} (x_i - \overline{X}) = 0$

3 Marks Questions

1. Find the median of the following data 19, 25, 59, 48, 35, 31, 30, 32, 51. If 25 is replaced by 52, what will be the new median.

Ans: 32, 35

i=1

2. If the mean of the following distribution is 6, then find the value of **p**.



X	2	4	6	10	p+5
f	3	2	3	1	2

Ans: 7

3. If the mean of five observations x, x+2, x+4, x+6, x+8 is 11, find the mean of first three observation.

Ans: 9

4. The mean of 5 numbers is 18. If one number is excluded, their mean is 16, find the excluded number.

Ans: 26

5. The following observations have been arranged in ascending order. If the median of the data is 63, find the value of x.

\$29,32,48,50,x,x + 2,72,78,84,95\$

Ans: 62

6. (i) Find the mean of the following data: 25, 27, 19, 29, 21, 23, 25, 30, 28, 20.

Show that the sum of deviation of all the observations from the mean is zero.

(ii) Find the median of the data given above.

Ans: (i) Mean = $\overline{x} = \frac{25 + 27 + 19 + 29 + 21 + 23 + 25 + 30 + 28 + 20}{10} = \frac{247}{10} = 24.7$

Sum of the deviations of all the observations from the mean $=\sum_{i=1}^{10} (x-\overline{x})$



$$=(25-24.7)+(27-24.7)+(19-24.7)+(27-24.7)+$$

$$(23-24.7)+(25-24.7)+(30-24.7)+(28-24.7)+(20-24.7)$$

= 0.3 + 2.3 - 5.7 + 4.3 - 2.7 - 1.7 + 03 + 5.3 + 3.3 - 4.7 = 0

(ii) Arranging the data in ascending order: \$19,20,21,23,25,25,27,28,29,30\$

$$\therefore \text{ Median} = \frac{\left[\frac{n}{2}\right]^{th} \text{ observation } + \left[\frac{n}{2} + 1\right]^{th} \text{ observation}}{2}$$

$$=\frac{5^{\text{th}} \text{ observation } + 6^{\text{th}} \text{ observation}}{2} = \frac{25 + 25}{2} = 25$$

- 7. If the mean of the following data is 21 . Find the value of P.
- X 7 15 28 5 1
- Y 8 20 P 3 2

Ans:

X _i	f_i	$f_1 \times x_i$
7	8	56
15	20	300
28	р	28p



5	3	15	
1	2	2	
Total	33+p	373+28p	
$\sum f_i =$	33+ <i>P</i>	$\sum x_i f_i = 37$	3+28P

$$\overline{x} = \frac{\sum x_1 f_f}{\sum f_i}$$

 $21 = \frac{373 + 28P}{33 + P}$

 $21 \times 33 + 21P = 373 + 28P$

693 + 21P = 373 + 28P

693 - 373 = 28P - 21P

320 = 7P

$$P = \frac{320}{7}$$

 $\therefore P = 45.7$

8. In a mathematics test given to 10 students, the following marks [out of 100] are recorded as: 82, 41, 39, 52, 53, 45, 96, 47, 50, 60.

Find out the mean & median of the above marks.

Ans: The given observation are: 82, 41, 39, 52, 53, 45, 96, 47, 50, 60

Sum of 10 observations = 82 + 41 + 39 + 52 + 53 + 45 + 96 + 47 + 50 + 60 = 565

Class IX Maths



$$\overline{X} = \frac{1}{n} \left[\sum_{i=1}^{n} x_i \right]$$

$$\therefore \bar{X} = \frac{565}{10} = 56.5$$

Arrange given observation in ascending order:

39, 41, 45, 47, 50, 52, 53, 60, 82, 96.

Here, n = 10 [even no.]

:. Median =
$$\frac{\left(\frac{n}{2}\right)^{\text{th}} \text{ observation } + \left(\frac{n}{2} + 1\right)^{\text{th}} \text{ observation}}{2}$$

$$=\frac{5^{\text{th}} \text{ observation } + 6^{\text{th}} \text{ observation}}{2}$$

$$=\frac{50+52}{2}=\frac{102}{2}=51$$

9. The following is the monthly expenditure (Rs.) of ten families of the particular area:

145, 115, 129, 135, 139, 158, 170, 175, 188, 163

a. Make a frequency distribution table by using the following class interval:

100-120, 120-140, 140-160, 160-180, 180-200.

(b) Construct a frequency polygon for the above frequency distribution.

Ans:

Frequency distribution



Class Intervals	Tally Marks	Frequency	
100-120	I	1	
120-140	III	3	
140-160	Ш	2	
160-180	III	3	
180-120	I	1	
Total		10	
Y			





10. The mean of 5 numbers is 39. If one number is excluded, their mean is 35, find the excluded number.

Ans: The mean of 5 numbers = 39

So, sum of five numbers

The mean of 4 numbers

So, sum of four numbers $= 35 \times 4 = 140$

Thus, Excluded numbers = Sum of five numbers - Sum of four numbers

=195 - 140 = 55

11. If the mean of 8 observation x, x+1, x+3, x+4, x+5, x+6, x+7 is 50, find the mean of first 5 observation.

Ans: Mean $=\frac{\sum_{n} x_{i}}{n}$ $\overline{x} = \frac{x + (x+1) + (x+2) + (x+3) + (x+4) + (x+5) + (x+6) + (x+7)}{8}$ $50 = \frac{8x + 28}{8}$ 400 - 28 = 8x $\therefore x = \frac{372}{8} = 46.5$ The given set of 8 observations is 46.5, 47.5, 48.5, 50.5, 49.5, 51.5, 52.5, 53.5.

So, mean of first 5 observations is:

$$\overline{x} = \frac{46.5 + 47.5 + 48.5 + 49.5 + 50.5}{5} = \frac{242.5}{5} = 48.5$$

12. Represent the following data by means of histogram.

Class IX Maths





Following will be the adjusted frequencies:

Weekly wages (in rupees)	Frequency	Adjusted frequencies
10-15	7	$\frac{5}{5} \times 7 = 7$
15-20	9	$\frac{5}{5} \times 9 = 9$



20-25	8	$\frac{5}{5} \times 8 = 8$
25-30	5	$\frac{5}{5} \times 5 = 5$
30-40	12	$\frac{5}{10} \times 12 = 6$
40-60	12	$\frac{5}{20} \times 12 = 3$
60-80	8	$\frac{5}{20} \times 8 = 2$

13. The Mean of monthly salary of 10 members of a group is Rs 1445, one more member whose monthly salary is Rs 1500 has joined the group. Find mean monthly salary of 11 member of the group.

Ans: Here n = 10, $\overline{X} = 1445$

So,
$$\overline{X} = \frac{1}{n} \left(\sum x_i \right) \Rightarrow n\overline{X} = \sum x_i \Rightarrow 10 \times 1445 = \sum x_i$$

$$\Rightarrow \sum x_i = 14450$$

So, total monthly wages of 10 person $R_s = 14450$

Monthly salary wages of one more person who joined the group is 1500.

 \therefore Total monthly wages of 11 persons = Rs 14450+1500 = Rs 15950

So, average monthly salary to 11 person



$$=\frac{\text{total monthly wages}}{11} = \frac{15950}{11} = \text{Rs}1450$$

14. The sum of the deviations of a set of n values x_1, x_2, \dots, x_n measured from 50 is -10 and the sum of deviation of the values from 46 is 70. Find the values of n and the mean.

Ans: We have,

$$\sum_{i=1}^{n} (x_i - 50) = -10 \text{ and } \sum_{i=1}^{n} (x_i - 46) = 70$$
$$\Rightarrow \sum_{i=1}^{n} x_i - 50n = 10 \rightarrow \text{ (i)}$$
$$\Rightarrow \sum_{i=1}^{n} x_i - 46n = 70 \rightarrow \text{ (ii)}$$

Subtracting (ii) from (i), we get -4n = -80

n = 20

Putting n = 20 in (i), we get

$$\sum_{i=1}^{n} x_i - 50 \times 20 = -10$$

$$\Rightarrow \sum_{i=1}^{N} X_i = 990$$

:. mean
$$=\frac{1}{n} \left[\sum_{i=1}^{n} x_i \right] = \frac{990}{20} = 49.5$$

So, n = 20 and mean = 49.5

15. There are 50 students in a class out of which 40 are boys and rest girls. The average weight of the class is 44 kg and the average weight of the girls is40 kg. Find the average weight of the boys.

Class IX Maths



Ans: We have, n = No. of students in a class = 50

 $n_1 = No. of boys in a class = 40$

 $n_2 = No. of girls in a class = 10$

 \overline{X}_1 = Average weight of boys =?

 X_2 = Average weight of girls = 40 kg

 $\therefore \overline{X} = \frac{n_1 \overline{X}_1 + n_2 \overline{X}_2}{n_1 + n_2}$ $\Rightarrow 44 = \frac{40 \overline{X}_1 + 10 \times 40}{40 + 10}$ $\Rightarrow 50 \times 44 = 40 \overline{X}_1 + 400$ $\Rightarrow 2200 = 40 \overline{X}_1 + 400$ $\Rightarrow \overline{X}_1 = 45$

Hence, the average weight of boys is 45Kg

16. The mean of 100 items was found to be 300. If at the time of calculation two itemswere wrongly taken as 32 and 12 instead of 23 and 11, find the correct mean.

Ans: Here, n = 100, X = 30

So,
$$\overline{X} = \frac{1}{n} \left(\sum x_i \right) \Rightarrow \sum x_i = n \overline{X}$$

$$\Rightarrow \sum x_i = 100 \times 30 = 3000$$

 \therefore Incorrect value of $\sum x_i = 3000$

Now, correct value of $\sum x_i$ = incorrect value of $\sum x_i - ($ sum of incorrect value) + (sum of correct value)

Class IX Maths



=3000 - (32 + 12) + (23 + 11) = 2990

Correct mean = $\frac{\text{Correct value of } \sum x_i}{n} = \frac{2990}{100} = 29.9$

17. The mean of 10 numbers is 20. If 8 is subtracted from every number, what will be the new mean?

Ans: Let x_1, x_2, \dots, x_{10} be 10 numbers with their mean equal to 20.

Then,

$$\overline{X} = \frac{1}{n} \left(\sum x_i \right)$$

$$20 = \frac{x_1 + x_2 + \dots + x_{20}}{10}$$

 $\Rightarrow 200 = \mathbf{x}_1 + \mathbf{x}_2 + \ldots + \mathbf{x}_{10}$

New numbers are $x_1 - 8, x_2 - 8, \dots, x_{10} - 8$

Let \overline{X} be the mean of new number.

$$\overline{X} = \frac{(x_1 - 8) + (x_2 - 8) + \dots + (x_{10} - 8)}{10}$$
$$X = \frac{(x_1 + x_2 + \dots + x_{10}) - 8 \times 10}{10}$$
$$= \frac{200 - 80}{10} = 12$$

18. The mean of n observation x_1, x_2, \dots, x_n , is \overline{X} . If (a-b) is added to each of the observation, show that the mean of the new set of observation is $\overline{X} + (a-b)$.

Ans: We have, $\overline{X} = \frac{x_1 + x_1 + \dots + x_n}{n} \rightarrow (i)$



Let X be the mean of
$$x_1 + (a-b), x_2 + (a-b), \dots, x_n + (a-b)$$
. Then,

$$X = \frac{[x_1 + (a-b)] + [x_2 + (a-b)] + \dots + [x_n + (a-b)]}{n}$$

$$= \frac{x_1 + x_2 + \dots + x_n + n(a-b)}{n}$$

$$= \frac{x_1 + x_2 + \dots + x_n}{n} + \frac{n(a-b)}{n}$$

$$= X + (a-b) \text{ [using (i)]}$$

19. If x_1, x_2, \dots, x_n are n values of a variable X such that $\sum_{i=1}^n (x_i - 2) = 110$ and

$$\sum_{i=1}^{n} (x_{1} - 5).$$
 Find the value of n and the mean.
Ans: We have:
$$\sum_{i=1}^{n} (x_{i} - 2) = 110 \text{ and } \sum_{i=1}^{n} (x_{i} - 5) = 20$$

$$\Rightarrow (x_{1} - 2) + (x_{2} - 2) + \dots + (x_{n} - 2) = 110$$

$$\Rightarrow (x_{1} - 5) + (x_{2} - 5) + \dots + (x_{n} - 5) = 20$$

$$\Rightarrow (x_{1} + x_{2} + \dots + x_{n}) - 2n = 110 \text{ and } (x_{1} + x_{2} + \dots + x_{1}) - 5n = 20$$

$$\Rightarrow \sum_{i=1}^{n} x_{i} - 2n = 110 \text{ and } \sum_{i=1}^{e} x_{1} - 5n = 20$$
Thus, we have $S - 2n = 110 \dots(i)$

and S - 5n = 20....(ii)

Subtracting (ii) from (i), we get:

$$3n = 90$$

$$n = 30$$



Putting n = 30 in (i), we get:

$$S-60 = 110 \implies S = 170 \implies \sum_{i=1}^{n} x_i = 170$$
$$\therefore \text{ mean } = \frac{1}{n} \left[\sum_{i=1}^{n} x_i \right]$$
$$= \frac{170}{30} = \frac{17}{3}$$

20. Find the missing frequencies in the following frequency distribution shown that the mean of the distribution is 1.46.

No. of accidents(x)		0	1	2	3	4	5	Total
Frequency(f)		45	?	?	25	10	5	200

Ans: Let the missing frequencies be f_1 and f_2 .

X _i	f_i	$f_i x_i$	
0	46	0	
1	f_1	f_1	
2	f_2	2 <i>f</i> ₂	
3	25	75	



4	10	40
5	5	25
	$\sum f_i = 86 + f_1 + f_2$	$\sum f_i x_i = 140 + f_1 + 2f_2$

We have: N = 200

- $\therefore 200 = 86 + f_1 + f_2$
- $114 = f_1 + f_2$(i)

Also, Mean =1.46

$$\Rightarrow 1.46 = \frac{\sum f_i x_i}{N}$$

$$\Rightarrow 1.46 = \frac{140 + f_1 + 2f_2}{200}$$

 $\Rightarrow 292 = 140 + f_1 + 2f_2$

 $\Rightarrow 152 = f_1 + 2f_2 \dots \dots (ii)$

Solving (i) and (ii), we get

$$f_1 = 76$$
 and $f_2 = 38$

21. Give some examples of data that you can collect from your day to day life.

Ans: Some examples of data that we can gather from our day to day life are:

• Number of students in our college.



- Number of fans in our school library.
- Electricity bills of our shop for last three years.
- Election results obtained from television or newspapers.
- Literacy rate figures obtained from National Survey.
- Heights of 30 students of our class.
- Maximum temperatures of the days of a particular week from weather forcast.
- Number of members in the families of your locality from a record.
- Distances from the college of the homes of five students.

22. Classify the data in Q21 above as primary or secondary data.

Ans: Primary data: When investigator himself collects the information with some specific objective in mind, data obtained is called Primary data. Such types of data are collected for first time for their own use. Above mentioned (i), (ii), (iii), (iv) and (ix) are the examples of primary data.

Secondary data: It includes the collection of information from sources which already had the information stored such as newspaper, television, etc. Above mentioned (iv), (v), (vii) and (viii) are the examples of secondary data.

23. The blood groups of 30 students of a class VIII are recorded as follows:

A, B, O, O, AB, O, A, O, B, A, O, B, A, O, B, A, O, O, A, AB, O, A, A, O, O, AB, B, A, O, B, A, B, O.

Represent this data in the form of a frequency distribution table. Which is the most common and which is the rarest blood group among these students?

Ans: The frequency distribution table for the given data is as follows:

V	ed	ar	t	il
LIVE	ONLINE	тит	ORIN	IG

Blood group	Tally Marks	Frequency (Number of students)
А	.¥Í Ⅲ	9
В	JHÍ I	6
0	JHT JHT II	12
AB	=	3
Total		30

From the table, It can be observed that the most common blood group is O and the rarest group is AB.

24. Distance (in km) of 40 engineers from their place of residence to their place of work were found as follows:

5	3	10	20	25	11	13	7	12	31
19	10	12	17	18	11	32	17	16	2
7	9	7	8	3	5	12	15	18	3
12	14	2	9	6	15	15	7	6	12



Construct a grouped frequency distribution table with class size 5 for the data given 40 above taking the first interval as 0 - 5 (5 not included). What main features do you observe from this tabular representation?

Ans. The grouped frequency distribution table for the given data is as follows:

Blood group	Tally Marks	Frequency (Number of female engineer's)
0-5	Ж(5
5-10		11
10-15		11
15-20		9
20-25		1
25-30	1	1
30-35	1	2
Total		40



From the table we observe that out of 40 female engineers 36(5+11+11+9) engineers i.e. 90% of the total female engineers reside less than 20 km from their place of work.

25. The relative hu	midity (in %	6) of a certain	city for a m	onth of 30	days was
as follows:					

98.1	98.6	98.2	98.3	86.5	95.3	92.9	<mark>9</mark> 6.3	94.2	95.1
89.2	92.3	97.1	93.5	92.7	95.1	97.2	93.3	95.2	97.3
96.2	92.1	84.9	90.2	95.7	98.3	97.3	96.1	92.1	89

(i) Construct a grouped frequency distribution table with classes 84-86,86-88 etc.

(ii) Which month or season do you think this data is about?

(iii) What is the range of this data?

Ans: (i) The grouped frequency distribution table for the given data is as follows:

Relative humidity (in %)	Tally Marks	No. o <mark>f days</mark>
84-86	_	1
86-88		1
88-90		2
90-92		2

Class IX Maths



92-94	II JK	7
94-96	_	6
96-98	II M	7
98-100		4

(ii) Data appears to be taken in the rainy season because we observe that relative humidity is high.

(iii) From the data, we observe

Highest relative humidity = 99.2%

Lowest relative humidity = 84.9%

Range = (99.2 - 84.9)% = 14.3%

26. The heights of 50 students, measured to the nearest centimeters have been found to be as follows:

161 150 154 165 168 161 154 162 150 1⁵1

162 164 171 165 158 154 156 172 160 170

153 159 161 170 1<mark>62 165 166 168 165 164</mark>

154 152 153 156 158 162 160 161 173 166

161 159 162 167 168 159 158 153 154 159

(i) Represent the data given above by a grouped frequency distribution table, taking the class - intervals as 160–165,165–170 etc.



(ii) What can you conclude about their heights from the table?

Ans: (i) The grouped frequency distribution table for the given data is as follows:

Distance	Tally Marks	No. of students Frequency
150-155	JM JM	12
155-160	X III	9
160-165	, WA (IIII	14
165-170	JHI JHI	10
170-175	Ш	5
Total		50

(ii) From the frequency distribution table drawn above, we conclude that more than 50% of the students are shorter than 165 cm.

27. A study was conducted to find out the concentration of sulphur dioxide in the air in parts per million (ppm) of a certain city. The data obtained for 30 days is as follows:



0.030.080.080.090.040.170.160.050.020.060.180.200.110.080.120.130.220.070.080.010.100.060.090.180.110.070.050.070.010.04

(i) Make a grouped frequency distribution table for this data with class intervals as

0.01 - 0.04, 0.04-0.08 and so on.

(ii) For how many days, was the concentration of sulphur dioxide more than 0.11 parts per million.

Ans: (i) The minimum and maximum concentration of Sulphur dioxide in the air in parts per million is 0.01 and 0.22 respectively.

It is given that 0.00-0.04 is one of the class intervals and the class size is the same.

So, the classes of equal size are

0.00-0.04, 0.04-0.08, ..., 0.20-0.24

Thus, the frequency distribution table is as under.

Concentration of Sulphur dioxide (in ppm)	Frequency
0.00 - 0.04	4
0.04 - 0.08	9
0.08 - 0.12	9
0.12-0.16	2
0.16-0.20	4
0.20-0.24	2
Total	30

(ii) The concentration of sulphur dioxide was more than 0.11 ppm for 8 days.



28. Three coins were tossed 30 times simultaneously. Each time the number of heads occurring was noted down as follows:

0 1 2 2 1 2 3 1 3 0 1 3 1 1 2 2 0 1 2 1 **Prepare a frequency distribution for the data** 3 0 0 1 1 2 3 2 0 **given above.**

Ans:

Number of heads	Num <mark>ber of times (frequency)</mark>
0	6
1	10
2	9
3	5
Total	30

29. The value of π up to 50 decimal places is given below:

3.1415926535 8979323846 2643383279 5028841971 6939937510

(a) Make a frequency distribution of the digits after the decimal point list the digits from 0 to 9 in your first column.

(b) What are the most and the least frequency occurring digits?

Ans: (i) The frequency distribution table is as under:



Digits	Frequency
0	2
1	5
2	5
3	8
4	4
5	5
6	4
7	4
8	5
9	8
Total	50

(ii) The most frequently occurring digits are 3 and 9.

The least occurring is 0.

30. Thirty children were asked about the number of hours they watched TV programmers in the previous week. The results were found as follows:

1	6	2	3	5	12	5	8	4	8
10	3	4	12	2	8	15	1	17	6
3	2	8	5	9	6	8	7	14	12

i. Make a grouped frequency distribution table for this data, taking class width 5 and one of the class interval 5 – 10.

ii. How many children watched television for 15 or more hours a week?

Ans: (i) The minimum and maximum number of hours children watched TV programmers in the previous week are 1 hour and 17 hours, respectively. It is given that 5 - 10 is one of the class intervals and the class size is same. So, the classes of equal size are 0 - 5, 5 - 10, 10 - 15, 15 - 20.

Thus, the frequency distribution table is as under.



Number of heads	Frequency	
0-5	10	
5-10	13	
10-15	5	
15 - 20	2	
Total	30	

(ii) 2 children watched television for 15 or more hours a week

31. A company manufactures car-batteries of particular type. The live (in years) of 40 such batteries were recorded as follows:

2.6	3.0	3.7	3.2	<mark>2</mark> .2	4.1	3 <mark>.5</mark>	4.5
3.5	2.3	3.2	3.4	<u>3.8</u>	3.2	4. <mark>6</mark>	3.7
2.5	4.4	3.4	3.3	2.9	3.0	4.3	2.8
3.5	3.2	3.9	3.2	3.2	3.1	3.7	3.4
4.6	3.8	3.2	2.6	3.5	4.2	2.9	3.6

Construct a grouped frequency distribution table for this data, using class intervals of 45 size 0.5 starting from the interval 2 - 2.5.

Ans: The minimum and maximum life (in number of years) of car batteries are 2.2 years and 4.6 years. It is given that 2 - 2.5 is one of the class interval with uniform size of 0.5. So, the classes of equal size are 2.0 - 2.5, 2.5 - 3.0, 3.0 - 3.5,...., 4.5 - 5.0.

Thus, the frequency distribution table is as under:



Frequency
2
6
14
11
4
3
30

4 Marks Questions

1. Find the value of x and y in following distribution if it known that the mean of the distribution is 1.46.

Ans: x = 76, y = 38

2. The mean monthly salary of 10 members of a group is Rs 1445, one more member whose monthly salary is Rs 1500 has joined the group. Find the mean monthly salary of 11 members of the group.

Ans: Rs 1450

3. Given below is a cumulative frequency distribution table showing the age of people living in a locality.



Age in years	No. of persons
Above 108	0
Above 96	1
Above 84	3
Above 72	5
Above 60	20
Above 48	1 <mark>58</mark>
Above 36	42 <mark>7</mark>
Above 24	809
Above 12	1026
Above 0	1124

Prepare a frequency distribution table.

Ans:

Marks	0-12	12-24	24-36	<mark>36–48</mark>	48-60	<mark>6</mark> 0-72	72 <mark>-84</mark>	84-96	96-108
Person	98	217	382	269	138	15	2	2	1

4. If $\mathbf{x}, \mathbf{x}_2, \dots, \mathbf{x}_n$ are n values of a variable x such that $\sum_{i=1}^n (x_i - 2) = 110$ and $\sum_{i=1}^n (x_i - 5) = 20$ find the value of n and mean.

Ans: n = 30, mean = $\frac{17}{3}$

5. The mean of 200 items was 50. Later on, it was discovered that the two items were misread as 92 and 8 instead of 192 and 88. Find the correct mean.

Ans: 50.9

6. Find the value of p, if the mean of following distribution is 20.



x 15 17 19 20+p 23 frequency 2 3 4 5p 6

Ans: 1

7. Prove that the sum of the deviations of individual's observations from the mean is zero.

Ans: Let $x_1, x_2, x_3, \dots, x_n$ be n individual observations whose mean is \overline{x} . The sum of the deviations of these n observations from \overline{x} is given by:

$$\sum_{i=1}^{n} (x_i - \overline{x}) = (x_i - \overline{x}) + (x_2 - \overline{x}) + \dots + (x_n - \overline{x})$$

Also, we have $\overline{x} = \frac{1}{n} \left[\sum_{i=1}^{n} x_i \right]$
 $\Rightarrow n\overline{x} = \sum_{i=1}^{n} x_i$
Now, $\sum_{i=1}^{n} (x_i - \overline{x}) = (x_i - \overline{x}) + (x_2 - \overline{x}) + \dots + (x_n - \overline{x})$
 $= (x_1 + x_2 + \dots + x_n) - n\overline{x}$
 $= \sum_{i=1}^{n} x_i - n\overline{x}$
 $= n\overline{x} - n\overline{x}$
 $= 0$
So, $\sum_{i=1}^{n} (x_i - \overline{x}) = 0$

8. From the data given below find:



- (a) Lower limit of the third class
- (b) Upper limit of the seventh class.
- (c) Class boundaries of the sixth class.
- (d) The class mark of the fifth class.
- (e) The size of the second class
- (f) Draw histogram of the data.

No.ofWage[Rs.]150-200200-250250-300300-350350-400400-450450-500Employees5356875

Ans: (a) Lower limit of the third class = 250

- (b) Upper limit of the seventh class = 500
- (c) Class boundaries of the sixth class = 400 450
- (d) The class mark of the fifth class = 375
- (e) The size of the second class = 50
- (f) Histogram



9. Draw a histogram with frequency polygon for the following data.

class interval	25 - 29	30 - 34	<mark>35 –</mark> 39	40 - 44	45-49	50 - 54
freq <mark>uency</mark>	5	15	23	20	10	7

Ans: Ascertainment of lower and upper class limits: since the difference between the second and first mid-points is 25-29

Let h = 1

Then for continuous frequency distribution, we subtract $\frac{h}{2}$ from lower limit and Add $\frac{h}{2}$ to upper limit.





Ans: Let the number of girls and boys be n_1 and n_2 respectively.

We have:



- $\overline{X_1}$: Average score of girls = 67
- $\overline{X_2}$: Average score of boys = 63
- *X* : Average score of the whole class = 64.5

$$X = \frac{n_1 \overline{X_1} + n_2 \overline{X_2}}{n_2 + n_2}$$
$$\Rightarrow 64.5 = \frac{67n_1 + 63n_2}{n_1 + n_2}$$

- $\Rightarrow 64.5n_1 + 64.5n_2 = 67n_1 + 63n_2$
- \Rightarrow 2.5n₁ = 1.5n₂
- $\Rightarrow 25n_1 = 15a_2$
- \Rightarrow 5n₁ = 3n₂

Total number of students in the class $= n_1 + n_2$

: percentage of girls $=\frac{n_1}{n_1+n_2} \times 100$

$$= \frac{n_1}{n_1 + \frac{5n_1}{3}} \times 100$$
$$= \frac{3n_1}{3n_1 + 5n_1} \times 100$$

$$=\frac{3}{8}\times100=37.5$$

Percentage of boys = $\frac{n_2}{n_1 + n_2} \times 100$

$$=\frac{n_2}{\frac{3n_2}{5}+n_2} \times 100$$



 $=\frac{5n_2}{3n_2+5n_2}\times 100$

= 62.5

11. Find the unknown entries (a, b, c, d, e, f) from the following frequency distribution of heights of 50 students in a class.

Class Intervals (height in cm)	Frequency	Cumulative frequency
150-155	12	a
155-160	b	25
160-165	10	с
165-170	d	43
170-175	e	48
175-180	2	f

Ans: Since the given frequency distribution is the frequency distribution of 50 students.

Therefore, g = 50

From the table, we have

a=12, b+12=25, 12+b+10=c, 12+b+10+d=43

12+b+10+d+e=48 and 12+b+10+d+e+g=f

Now,

 $b+12=25 \Longrightarrow b=13$



12+b+10=c $\Rightarrow 12+13+10=c$ $\Rightarrow c=35$ 12+b+10+d-43 $\Rightarrow 12+13+10+d=43$ $\Rightarrow d=8$ 12+b+10+d+e=48 $\Rightarrow 12+13+10+8+e=48$ $\Rightarrow e=5$ 12+b+10+d+e+2=f $\Rightarrow 12+13+10+8+5+2=f$ $\Rightarrow f=50$